




<b>SPECIES</b>	<b><i>Iva axillaris</i> Pursh ssp. <i>robustior</i> (Hook.) Bassett</b>	
<b>NRCS CODE: IVAXR</b>	<p>Family: <b>Asteraceae</b>  Order: <b>Asterales</b>  Subclass: <b>Asteridae</b>  Tribe: <b>Helenieae</b>  Subtribe: <b>Ambrosiinae</b>  Class: <b>Magnoliopsida</b></p>  	<p>A. Montalvo, w. Riverside Co.</p>
<b>Subspecific taxa</b>	California plants have been treated as a subspecific taxon of <i>I. axillaris</i> Pursh.	
<b>Synonyms</b>	<i>I. axillaris</i> Pursh var. <i>pubescens</i> Gray	
<b>Common name</b>	povertyweed, deathweed, devil's-weed, poverty sumpweed, bozzleweed, salt sage, marsh elder, small-flowered marsh elder (DiTomaso & Healy 2007, Painter 2009). There are many more common names.	
<b>Taxonomic relationships</b>	Species in the genus <i>Iva</i> are related to <i>Ambrosia</i> and <i>Artemisia</i> in southern California. Seven <i>Iva</i> species are currently recognized in North America (FNA), but nine species were assessed for phylogenetic relationships using cpDNA (Miao et al. 1995).	
<b>Related taxa in region</b>	<i>Iva hayesiana</i> A. Gray (rare, native subshrub found mostly in sw San Diego County) is the closest relative in the region. The species differ in chromosome number, stature, size and habitat; <i>I. hayesiana</i> has n=17 chromosomes, is a coastal (Miao et al. 1995), and also differs in having flower heads with separate phyllaries (Munz 1974, DiTomaso & Healy 2007). Ragweeds are in the same section of the Asteraceae. Some (e.g., <i>Ambrosia psilostachya</i> ) are found in similar areas and have a similar growth habit, but unlike <i>Iva</i> , they have incised or lobed leaf margins (DiTomaso & Healy 2007). Outside focus area: <i>Iva nevadensis</i> Jones is an annual herb found in sagebrush scrub and pinyon-juniper woodlands of CA (Munz 1974); <i>Iva acerosa</i> (Nutt.) Jackson is a perennial found on alkali sinks and desert areas of CA (Munz 1974).	
<b>Taxonomic issues</b>	Munz (1974), Hickman (1993) and JepsonOnline (1st edition, 2010) consider southern CA forms to be <i>I. axillaris</i> ssp. <i>robustior</i> . However, the JepsonOnline 2nd Ed (2010) may dismiss the varietal status. Subspecies <i>robustior</i> and <i>axillaris</i> can be separated, in part, by the degree of fusion of the outer phyllaries, leaf shape, and color of glands after drying (Best 1975). In ours, the fused phyllaries form a distinct cup. FNA (2010) and USDA PLANTS (2010) do not recognize subspecific taxa.	
<b>Other</b>	In some species, such as <i>Phragmites australis</i> (Saltonstall 2002, Vasquez et al. 2005), there are invasive and non-invasive genotypes; non-native aggressive genotypes from Eurasia have higher salt tolerance and have invaded North American wetlands, displacing many species over sometimes large areas. Such examples underline the value of using native non-invasive genotypes of taxa that are known to be invasive in other areas. Studies are needed to determine if there are differences in invasive potential within <i>I. axillaris</i> , especially between southwestern and northern subtaxa. This species is listed as a California Noxious Weed (with respect to agriculture), category CW (USDA PLANTS 2010, GRIN); however, its weediness is more of a problem in Canada and outside California (Munz & Keck 1968). The ssp. <i>robustior</i> is native to CA, the western US and western Canada and is not on the US Federal noxious weed list. However, the species is considered a noxious weed and serious pest in the prairies of Canada (Best 1975) and where it has been introduced in Australia (Parsons & Cuthbertson 1992). To avoid invasiveness issues, do not use around sensitive plant species or agricultural fields. Avoid introducing genotypes from areas where the plant has been introduced and is invasive and avoid placing seed increase plots in seed dispersal distance of other crops. Local seed laws may limit the sale and distribution of this species. Plants are somewhat resistant to 2-4-D at rates normally used for crops (Best 1975) and can be difficult to eradicate from agricultural fields.	

GENERAL	
<b>Map</b>	<p>Data provided by the participants of the Consortium of California Herbaria represent 144 records with coordinate data out of 330 records retrieved; data accessed 9/18/10. See Berkeley Mapper: <a href="https://ucjeps.berkeley.edu/consortium/">https://ucjeps.berkeley.edu/consortium/</a></p> 
<b>Geographic range</b>	<p><i>L. a. ssp. robustior</i>: Widespread; CA, British Columbia, MT, Central US, TX, and west of the Continental Divide (Bassett et al. 1962, Hickman 1993).  <i>L. a. ssp. axillaris</i> is widespread east of the Continental Divide in the northern US and Canada (Bassett et al. 1962).</p>
<b>Distribution in California; Ecological section and subsection</b>	<p>Common, but scattered throughout state (Munz &amp; Keck 1968, Hickman 1993). Many ecological sections (Goudey &amp; Smith 1994; Cleland et al. 2007), including: Modoc Plateau (M261G), Great Valley (262A), Central California Coast (261A), Sierra Nevada and Sierra Nevada Foothills (M261E,F), Mono and Southeastern Great Basin (341D,F), w Mojave Desert (322A), Southern California Mountains and Valleys (M262B), Southern California Coast (261B) .</p>
<b>Life history, life form</b>	<p>Long-lived perennial herb, capable of long periods of dormancy (Best 1975).</p>
<b>Distinguishing traits</b>	<p>Mat-forming perennial from vigorous creeping roots 0.2 to 0.6 dm tall; produces small nodding green flower heads from axils of leaves have outer phyllaries united into a cup; plants have simple, generally entire, pubescent, alternate leaves (1.5-2.5 cm long) and an unpleasant aromatic scent (Munz 1974, DiTomaso &amp; Healy 2007, FNA 2010). Plants can be slightly woody at base and leaves have glandular dots (Munz 1974).</p>
<b>Root system, rhizomes, stolons, etc.</b>	<p>Creeping, woody, highly branched root system (DiTomaso &amp; Healy 2007). Can spread vigorously from root system, potentially forming large clones (Best 1975).</p>
<b>Rooting depth</b>	<p>To 2.5 m depth (Best 1975, DiTomaso &amp; Healy 2007).</p>
HABITAT	
<b>Plant association groups</b>	<p>Coastal salt marsh and alkali plains communities (DiTomaso &amp; Healy 2007)</p>
<b>Habitat affinity and breadth of habitat</b>	<p>Weedy plant found from deserts to coastal areas in alkaline plains, depressions, or edges of saline marshes (Munz 1974); especially found in agricultural areas, pastures, rangeland, roadsides, and other disturbed areas (DiTomaso &amp; Healy 2007).</p>
<b>Elevation range</b>	<p>Sea level to 2500 m (Hickman 1993).</p>
<b>Soil texture, chemicals, depth</b>	<p>Found growing on poorly drained alkaline and saline soils, but also found on more favorable neutral, well draining soils (DiTomaso &amp; Healy 2007); in one study (primarily Canada) plants were found in soils with pH ranging from 6.59-8.77 and sodium content from 2.83 to 6.10 (Best 1975).</p>
<b>Drought tolerance</b>	<p>Plants typically grow in seasonally moist depressions, washes, along riparian edges and near streams or irrigation ditches. Plants can be found on dry slopes above riparian areas in Riverside Co., but it is possible that the deep root systems had contacted capillary water (A. Montalvo, pers obs).</p>
<b>Precipitation</b>	<p>Occurs under a broad range of precipitation normals, but in low precipitation regions, tends to occurs in areas that are seasonably wet (JepsonOnline 2010).</p>
<b>Flooding or high water tolerance</b>	<p>Occurs in seasonally wet, saline or alkaline areas (FNA 2010) and in areas that flood occasionally.</p>
<b>Wetland indicator status for California</b>	<p>Facultative (USDA PLANTS 2010).</p>
<b>Shade tolerance</b>	<p>Full sun to partial shade (A. Montalvo, pers. obs.)</p>

<b>GROWTH AND REPRODUCTION</b>	
<b>Seedling emergence relevant to general ecology</b>	Best (1957) reported that in natural ecosystems, there is generally very little seed germination, as most seeds are not viable due to seed predation. Most studies have been in Canada and the northern US.
<b>Growth pattern, phenology</b>	Flowers May-Sept. in California (Hickman 1993); foliage will die back in winter but roots remain alive (DiTomaso & Healy 2007). In North America, flowers May to October (FNA 2009).
<b>Vegetative propagation</b>	Clonal reproduction from creeping roots (DiTomaso & Healy 2007). Root pieces will lead to propagation and spread of plant (Best 1975, Parsons & Cuthbertson 1992).
<b>Regeneration after fire or other disturbance</b>	Investigation is needed to see if plants resprout after wildfire. Plants withstand mowing and other mechanical disturbance (DiTomaso & Healy 2007).
<b>Pollination</b>	Wind pollinated (Colin & Jones 1980, DiTomaso & Healy 2007).
<b>Primary seed dispersal</b>	The achenes are small and light, but they have no specialized dispersal structures. The seeds are also buoyant enough to be carried by water after they drop to the ground (Bruns & Rasmussen 1953).
<b>Breeding system, mating system</b>	Large clones can be expected to self-pollinate, but self-pollination resulted in underdeveloped embryos for five populations tested, including both subspecies (Bassett et al. 1962). This can be caused by self-incompatibility or severe inbreeding depression. This suggests seeds are produced by outcrossing and that multiple genotypes are necessary for seed production.
<b>Hybridization potential</b>	Potential for cross pollination of plants from different populations or varieties is high in this wind pollinated plant. In Colorado and Wyoming, ssp. <i>axillaris</i> and ssp. <i>robustior</i> overlap in distribution. Plants intermediate in form between have been observed, suggesting hybridization (Best 1975).
<b>Inbreeding and outbreeding effects</b>	Self-pollination resulted in underdeveloped embryos (Bassett et al. 1962). This may be an expression of inbreeding depression that occurs very early in embryo development. Self-incompatibility is also likely.
<b>BIOLOGICAL INTERACTIONS</b>	
<b>Competitiveness</b>	In agricultural areas, <i>I. axillaris</i> can significantly reduce crop yields in the U.S. and Canada (Best 1975, DiTomaso & Healy 2007). Extracts from leaves reduce germination and growth of roots and shoots in wheat, grasses, and legumes (Parsons & Cuthbertson 1992).
<b>Herbivory, seed predation, disease</b>	Seeds are eaten by larvae of the beetle <i>Smicronyx utilis</i> Buchanan (Best 1975). Another beetle, <i>Zygospila conjuncta conjuncta</i> , as well as other insects including <i>Rhopalus viridicatus</i> , <i>Psylliodes punctulata</i> , <i>Systema blada</i> , <i>Monoxia angularis</i> , <i>Collop vittatus</i> , and <i>Hippodamia parenthesis</i> are known to feed on different parts of the plant (Best 1975). <i>Ophaella communa</i> LaSage feeds on <i>Iva axillaris</i> in CA, and <i>I. frutescens</i> in eastern NA (Futuyma et al. 1993). In addition, researchers found that there was genetic variation in the insect with respect to larval survival on host plants within chemically similar plants in the Asteraceae. The greatest variation was with <i>I. frutescens</i> .
<b>Palatability, attractiveness to animals, response to grazing</b>	A weed database by the California Department of Food and Agriculture reported that plants are not palatable and are seldom eaten by livestock ( <a href="http://www.cdffa.ca.gov/phpps/ipc/weedinfo/iva-axillaris.htm">http://www.cdffa.ca.gov/phpps/ipc/weedinfo/iva-axillaris.htm</a> link no longer active March 2020); however, Hanley & Hanley (1982) reported that <i>I. axillaris</i> was grazed in the Great Basin by sheep. The plants have been reported to absorb selenium from the soil and accumulate sufficient quantities to be poisonous to some animals (Best 1975).
<b>Mycorrhizal?</b>	Found to be colonized with vesicular-arbuscular mycorrhizal fungi in disturbed areas of UT (Pendleton & Smith 1983).
<b>ECOLOGICAL GENETICS</b>	
<b>Ploidy</b>	2n=36, 54 (Hickman 1993). No correlates between variation in ploidy and morphology have been found (Best 1975). Miao et al. (1995) found <i>I. axillaris</i> to be the most primitive member of the genus based on cpDNA, chromosome number, and flower head structure. Primitive number of the tribe is n=18, and <i>Iva axillaris</i> ssp. <i>robustior</i> n=18 (Payne et al. 1964).
<b>Plasticity</b>	No information.
<b>Geographic variation (morphological and physiological traits)</b>	There is some morphological differentiation of populations classified as different subspecies, but no studies have been found that document the genetic basis of the variation.

<b>Genetic variation and population structure</b>	No studies found. Wind pollinated, self-incompatible plants typically have high rates of gene dispersal and low levels of population structure. Seed dispersal by water can also add to higher gene dispersal and low structure, but seeds dispersed by gravity could result in localized family structure. The clonal spread of these plants is likely to result in large patches of single genotypes.
<b>Phenotypic or genotypic variation in interactions with other organisms</b>	
<b>Local adaptation</b>	
<b>Translocation risks</b>	As mentioned above, the eastern genotypes may be more invasive than forms that occur in California. Care should be taken to avoid plants that are too aggressive.
<b>SEEDS</b>	For RSABG seed image see: <a href="http://www.hazmac.biz/USDA/USDAIvaAxillaris.html">http://www.hazmac.biz/USDA/USDAIvaAxillaris.html</a>
<b>General</b>	Seeds (achenes) are about 2 mm long and turnip-shaped (DiTomasso & Healy 2007).
<b>Seed longevity</b>	Seeds may be relatively short-lived both in the wild and in storage based on a review by Best (1975).
<b>Seed dormancy</b>	"Newly matured seed can be dormant" (DiTomaso & Healy 2007).
<b>Seed maturation</b>	
<b>Seed collecting</b>	This species may be similar to <i>Ambrosia</i> or <i>Artemisia</i> for which seeds can be hand-stripped or beaten into a hopper or open container (e.g., Stevens et al. 1996).
<b>Seed processing</b>	Wall and MacDonald (2009) recommend rubbing the flower material of similar <i>Artemisia</i> and <i>Ambrosia</i> over a sieve or medium screen, and then using a blower (with an Oregon Seed Blower unit) to remove the chaff.
<b>Seed storage</b>	Seeds may be relatively short-lived in dry storage. Seeds did not germinate when tested at any time during 22 and 60 months of dry storage (Bruns & Rasmussen 1953, 1957). However when stored submerged inside lumite screen bags in a Washington Canal, seeds germinated after prolonged storage (Bruns & Rasmussen 1953). There was no germination before 2 months, 43% germination after 7 months, < 3% germination after 10 months, increased to 38% after 14 months, then dropped again after 18 months. Germination cycled and increased with a decrease in water temperature. In another study, submerged seeds slowly germinated or disintegrated for 60 months, but 86% were no longer sound after 30 months (Bruns & Rasmussen 1957).
<b>Seed germination</b>	Scarification is not effective in promoting germination (Best 1975); seeds submerged outdoors in a canal in Washington had their highest germination at between months 6 to 8 months and again at 14 to 18 months when water temperatures were between 32 and 50 °F. Few seeds germinated when water temperatures were between 50 and 72 °F (Bruns & Rasmussen 1953). This suggests most seedling emergence will be in the cool season. Studies are needed to determine if cold stratification could aid germination of seeds from southern California plants.
<b>Seeds/lb</b>	161,997 (Stevens 1932).
<b>Planting</b>	
<b>Seed increase activities or potential?</b>	
<b>USES</b>	
<b>Revegetation and erosion control</b>	Promising for use on slopes of water quality basins and bioswales where alkalinity is expected. The root system of this species should be good for stabilizing salty or alkaline soils around water quality basins and seasonally wet bioswales.
<b>Habitat restoration</b>	Use cautiously and consider if too competitive for natural site.
<b>Horticulture or agriculture</b>	No information was found about how to grow plants. We only found information about how to get rid of <i>I. axillaris</i> from agricultural fields.
<b>Wildlife value</b>	In the mid-1900's in Oregon <i>I. axillaris</i> was found to make up more than 1% of the diet of antelopes (Mason 1952). There was also a low level of grazing in August by bison in Colorado (Peden 1976).
<b>Plant material releases by NRCS and cooperators</b>	None.

<b>Ethnobotanical</b>	Many uses by Native Americans have been reported in the ethnobotanical database ( <a href="http://naeb.brit.org/uses/species/2021/">http://naeb.brit.org/uses/species/2021/</a> ). For example, plants were used for contraception, as an abortifacient, gastrointestinal aid, and leaves were used as a plaster or in an infusion to treat sores or skin irritations. Pollen and hairs on leaves may cause an allergic reaction in sensitive individuals (DiTomaso & Healy 2007).
<b>ACKNOWLEDGMENTS</b>	Partial funding for production of this plant profile was provided by the U.S. Department of Agriculture, Forest Service, Pacific Southwest Region Native Plant Materials Program.
<b>CITATION</b>	Montalvo, A. M., L. K. Goode, and J. L. Beyers. 2010. Plant Profile for <i>Iva axillaris</i> . Native Plant Recommendations for Southern California Ecoregions. Riverside-Corona Resource Conservation District and U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Riverside, CA. Online: <a href="https://www.rcrd.org/plant-profiles">https://www.rcrd.org/plant-profiles</a> .
<b>LINKS TO REVIEWED DATABASES &amp; PLANT PROFILES</b> (last updated 3/24/2020)	
<b>Fire Effects Information System (FEIS)</b>	No matches: <a href="https://www.feis-crs.org/feis/">https://www.feis-crs.org/feis/</a>
<b>Jepson Flora, Herbarium (JepsonOnline)</b>	<a href="https://ucjeps.berkeley.edu/cgi-bin/get_cpn.pl?IVAX">https://ucjeps.berkeley.edu/cgi-bin/get_cpn.pl?IVAX</a>
<b>Jepson Flora, Herbarium, (JepsonOnline 2nd Ed.) updated to Jepson eFlora</b>	<a href="https://ucjeps.berkeley.edu/eflora/eflora_display.php?tid=3654">https://ucjeps.berkeley.edu/eflora/eflora_display.php?tid=3654</a>
<b>USDA PLANTS</b>	<a href="https://plants.usda.gov/core/profile?symbol=IVAX">https://plants.usda.gov/core/profile?symbol=IVAX</a>
<b>Native Plant Network Propagation Protocol Database (NPNDP)</b>	<a href="https://nnp.rngr.net/propagation/protocols">https://nnp.rngr.net/propagation/protocols</a>
<b>Native Seed Network</b>	<a href="https://nativeseednetwork.org/">https://nativeseednetwork.org/</a>
<b>GRIN</b>	<a href="https://npgsweb.ars-grin.gov/gringlobal/search.aspx?">https://npgsweb.ars-grin.gov/gringlobal/search.aspx?</a>
<b>Flora of North America (FNA) (online version)</b>	<a href="http://www.efloras.org/florataxon.aspx?flora_id=1&amp;taxon_id=250067018">http://www.efloras.org/florataxon.aspx?flora_id=1&amp;taxon_id=250067018</a>
<b>Native American Ethnobotany Database (NAE)</b>	<a href="http://naeb.brit.org/">http://naeb.brit.org/</a>
<b>Calflora</b>	<a href="https://www.calflora.org/">https://www.calflora.org/</a>
<b>Rancho Santa Ana Botanic Garden Seed Program, RSABG seed photos</b>	<a href="http://www.hazmac.biz/rsabghome.html">http://www.hazmac.biz/rsabghome.html</a>

## Bibliography for *Iva axillaris* ssp. *robustior*

- Bassett, I. J., G. A. Mulligan, and C. Frankton. 1962. Poverty weed, *Iva axillaris*, in Canada and the United States. *Canadian Journal of Botany* **40**:1243-1249.
- Best, K. F. 1975. Biology of Canadian weeds. 10. *Iva axillaris* Pursh. *Canadian Journal of Plant Science* **55**:293-301.
- Bruns, V. F., and L. W. Rasmussen. 1953. The effects of fresh water storage on the germination of certain weed seeds: I. White top, Russian knapweed, Canada thistle, morning glory, and poverty weed. *Weeds* **2**:138-147.
- Bruns, V. F., and L. W. Rasmussen. 1957. The effects of fresh water storage on the germination of certain weed seeds. II. White top, Russian knapweed, Canada thistle, morning glory, and poverty weed. *Weeds* **5**:20-24.
- Cleland, D. T. F., J. A. Freeouf, J. E. Keys, G.J. Nowacki, C. A. Carpenter, W. H. McNab. 2007. Ecological Subregions: Sections and Subsections for the conterminous United States. Gen. Tech. Report WO-76D [Map on CD-ROM] (A.M. Sloan, cartographer). U.S. Department of Agriculture, Forest Service, Washington, DC.
- Colin, L. J., and C. E. Jones. 1980. Pollen energetics and pollination modes. *American Journal of Botany* **67**:210-215.
- DiTomaso, J. M., and E. A. Healy. 2007. Weeds of California and Other Western States. Vol. 1. University of California, Agriculture and Natural Resources, Oakland, CA.
- Eckert, R. E., Jr., A. D. Bruner, G. J. Klomp, and F. F. Peterson. 1973. Mountain meadow improvement through seeding. *Journal of Range Management* **26**:200-203.
- FNA 2010. FNA Editorial Committee. 1993+. Flora of North America north of Mexico. 10+ volumes. New York and Oxford. Available online: [http://beta.floranorthamerica.org/Published\\_Volumes](http://beta.floranorthamerica.org/Published_Volumes). Last accessed August 2010.
- Futuyma, D. J., M. C. Keese, and S. J. Scheffer. 1993. Genetic constraints and the phylogeny of insect-plant associations: Responses of *Ophraella communa* (Coleoptera: Chrysomelidae) to host plants of its congeners. *Evolution* **47**:888-905.
- Goeden, R. D., and J. A. Teerink. 1993. Phytophagous insect faunas of *Dicoria canescens* and *Iva axillaris*, native relatives of ragweeds, *Ambrosia* spp., in southern California, with analyses of insect associates of Ambrosiinae. *Annals of the Entomological Society of America* **86**:37-50.
- Goudey, C. B., and D. W. Smith. 1994. Ecological units of California: Subsections (map). San Francisco, CA: U.S. Department of Agriculture, Forest Service. Pacific Southwest Region. Scale 1:1,000,000; colored.
- Hanley, T. A., and K. A. Hanley. 1982. Food resource partitioning by sympatric ungulates on Great Basin rangeland. *Journal of Range Management* **35**:152-158.
- Hazlett, D. L., and N. W. Sawyer. 1998. Distribution of alkaloid-rich plant species in shortgrass steppe vegetation. *Conservation Biology* **12**:1260-1268.
- Hickman, J. C., editor. 1993. The Jepson Manual: Higher Plants of California. University of California Press, Berkeley, CA.
- Hyder, D. N., R. E. Bement, E. E. Remmenga, and C. Terwilliger, Jr. 1966. Vegetation-soils and vegetation-grazing relations from frequency data. *Journal of Range Management* **19**:11-17.
- JepsonOnline. 2010. The Jepson Manual Higher Plants of California; Online Version with 2nd edition notes. <http://ucjeps.berkeley.edu/jepman.html>. (updated to Jepson eFlora: <https://ucjeps.berkeley.edu/eflora/>)

- JepsonOnline 2ndEd. 2010. B. G. Baldwin after Strother. 2011, in press. *Iva* in B. G. Baldwin et al. The Jepson Manual: Vascular Plants of California, 2nd Edition. University of California Press., Berkeley, CA. Online: <http://ucjeps.berkeley.edu/tjm2/review/treatments/compositae.html#3658>. Accessed: October 3, 2010. (Updated to: <https://ucjeps.berkeley.edu/eflora/>)
- Kelsey, F. D. 1989. Notes on the fungi of Helena, Mont. *The Journal of Mycology* **5**:80-82.
- MacCracken, J. G., and R. M. Hansen. 1982. Herbaceous vegetation of habitat used by blacktail jackrabbits and Nuttall cottontails in southeastern Idaho. *American Midland Naturalist* **107**:180-184.
- Miao, B., B. Turner, and T. Mabry. 1995. Molecular phylogeny of *Iva* (Asteraceae, Heliantheae) based on chloroplast DNA restriction site variation. *Plant Systematics and Evolution* **195**:1-12.
- Mason, E. 1952. Food habits and measurements of Hart Mountain antelope. *The Journal of Wildlife Management* **16**:387-389.
- McInnis, M. L., and M. Vavra. 1987. Dietary relationships among feral horses, cattle, and pronghorn in southeastern Oregon. *Journal of Range Management* **40**:60-66.
- Moir, W. H., and M. J. Trlica. 1976. Plant communities and vegetation pattern as affected by various treatments in shortgrass prairies of northeastern Colorado. *The Southwestern Naturalist* **21**:359-371.
- Munz, P. A. 1974. *A Flora of Southern California*. University of California Press, Berkeley, CA.
- Munz, P. A., and D. D. Keck. 1968. *A California Flora with Supplement*. University of California Press, Berkeley.
- Palmer, W. A., and R. D. Goeden. 1991. The host range of *Ophraella communa* Lesage (Coleoptera: Chrysomelidae). *The Coleopterists Bulletin* **45**:115-120.
- Painter, E. 2009. Common (vernacular) names applied to California vascular plants. *in*. University of California Herbarium. Available online: (2016 update: [https://ucjeps.berkeley.edu/cgi-bin/get\\_painter\\_common.pl?3654](https://ucjeps.berkeley.edu/cgi-bin/get_painter_common.pl?3654))
- Parsons, W. T., and E. G. Cuthbertson. 1992. *Noxious Weeds of Australia*. Inkata Press, Melbourne, Australia.
- Payne, W. W., P. H. Raven, and D. W. Kyhos. 1964. Chromosome numbers in Compositae. IV. Ambrosieae. *American Journal of Botany* **51**:419-424.
- Peden, D. G. 1976. Botanical composition of bison diets on shortgrass plains. *American Midland Naturalist* **96**:225-229.
- Pendleton, R. L., and B. N. Smith. 1983. Vesicular-arbuscular mycorrhizae of weedy and colonizer plant species at disturbed sites in Utah. *Oecologia* **59**:296-301.
- Piper, G. L., and B. L. Mulford. 1980. Life history observations on *Smicronyx commixtus* Dietz (Coleoptera: Curculionidae) in southeastern Washington. *The Coleopterists Bulletin* **34**:295-298.
- Saltonstall, K. 2002. Cryptic invasion by a non-native genotype of the common reed, *Phragmites australis*, into North America. *Proceedings of the National Academy of Sciences of the United States of America* **99**:2445-2449.
- Scheppegrell, W. 1917. Hay fever: Its cause and prevention in the Rocky Mountain and Pacific States. *Public Health Reports (1896-1970)* **32**:1135-1152.
- Standley, P. C. 1916. Fungi of New Mexico. *Mycologia* **8**:142-177.
- Stevens, O. A. 1932. The number and weight of seeds produced by weeds. *American Journal of Botany* **19**:784-794.
- Stevens, R., K. R. Jorgensen, S. A. Young, and S. B. Monsen. 1996. *Forb and shrub seed production guide for Utah*. Utah State University Extension, Logan, UT.

USDA PLANTS. 2010. The PLANTS Database (<https://plants.usda.gov/java/>). National Data Center, Baton Rouge, LA 70874-4490 USA.

Vasquez, E., E. Glenn, J. Brown, G. Guntenspergen, and S. Nelson. 2005. Salt tolerance underlies the cryptic invasion of North American salt marshes by an introduced haplotype of the common reed *Phragmites australis* (Poaceae). *Marine Ecology Progress Series* **298**:2005.

Wall, M., and J. Macdonald. 2009. Processing Seeds of California Native Plants for Conservation, Storage, and Restoration. Rancho Santa Ana Botanic Garden Seed Program, Claremont, CA; available online: <http://www.hazmac.biz/seedhome.html>.